

Estuarine caridean shrimps (Crustacea: Decapoda) from Ilhéus, Bahia, Brazil: Updated checklist and a key for their identification

Alexandre Oliveira Almeida*, Ana Carla Costa-Souza, Andressa Maria Cunha, Patricia Souza Santos, Mário Vitor Oliveira and Guidomar Oliveira Soledade

Universidade Estadual de Santa Cruz, Departamento de Ciências Biológicas. Rodovia Jorge Amado, km 16. 45662-900. Ilhéus, Bahia, BA, Brazil.

* Corresponding author. E-mail: aalmeida@uesc.br

ABSTRACT: We provide an updated list of the 22 species of caridean shrimps occurring in estuaries at Ilhéus, state of Bahia, Brazil, in the following families: Palaemonidae (4 species), Alpheidae (15 species), Hippolytidae (2 species) and Ogyrididae (1 species). The alpheid *Automate* cf. *dolichognatha* De Man, 1888 and the ogyridid *Ogyrides alphaerostris* (Kingsley, 1880) are reported from Bahia for the first time. The alpheids *Alpheus brasileiro* Anker, 2012, *A. buckupi* Almeida, Terossi, Araújo-Silva and Mantelatto, 2013, *A. chacei* Carvacho, 1979, *A. nuttingi* (Schmitt, 1924), *Leptalpheus axianassae* Dworschak and Coelho, 1999 and *Salmoneus carvachoi* Anker, 2007 are recorded from Ilhéus for the first time. *Alpheus angulosus* McClure, 2002 and *A. carlae* Anker, 2012 were previously reported from Ilhéus as *A. armillatus* (H. Milne Edwards, 1837). A key for identification of the carideans from estuaries of Ilhéus is provided.

INTRODUCTION

With almost 3500 species described so far, caridean shrimps are the second most diverse decapod infra-order and the most diverse group among the shrimp-like decapods (De Grave and Fransen 2011). Caridean shrimps are also remarkable for their ecological diversity, occurring from tropical to polar regions, in intertidal, subtidal and pelagic habitats, on hard and soft bottoms as epi- or infaunal organisms, on algae and seagrass, or in symbiosis with other animals (Bauer 2004). They have also successfully colonized diverse freshwater environments (De Grave *et al.* 2008), as well as estuaries and mangroves (*e.g.*, Carvacho 1979; Echeverría-Sáez 2003; Neves *et al.* 2007). Similarly to other decapods, caridean shrimps are permanent inhabitants of estuaries or use them as nursery grounds. As abundant members of the benthic communities in this type of environment, they are also an important link in aquatic food webs (Hooks *et al.* 1976; Gore *et al.* 1981; Eggleston *et al.* 1998; Neves *et al.* 2007).

Almeida *et al.* (2006), based on a survey conducted between 2003 and 2005, reported 51 species of decapod crustaceans in the estuaries of Ilhéus, state of Bahia, Brazil, of which 12 were caridean shrimps, belonging to three families (Palaemonidae, Alpheidae, and Hippolytidae). New samples were taken in 2011 and 2012, resulting in several new records for the study area. In Brazil, the caridean shrimps are less studied than other crustaceans such as members of the infraorders Brachyura and Anomura. This is due, in part, to the complicated taxonomy of certain groups, the lack of taxonomic revisions, and the absence of comprehensive keys for identification of Brazilian species. This contribution provides an updated list of estuarine caridean shrimps from Ilhéus. We also provide a key for identification of the species in the area.

MATERIALS AND METHODS

Study area

The Municipality of Ilhéus, located on the southeastern coast of the state of Bahia (Figure 1), northeastern Brazil, covers an area of 1,712 km². Its coastline is about 80 km long, is limited to the north by the Sargi River (14°30'06.7"S, 39°02'29.4"W) and to the south by the Acuípe River (15°05'41"S, 38°59'50"W), and includes several estuaries (Andrade 2003; Almeida *et al.* 2006). The Cachoeira River is the main river of Ilhéus, and together with the Santana and Fundão rivers forms a large estuary in the Ilhéus urban area. The Cachoeira River basin receives inputs from domestic and industrial effluents from Ilhéus and Itabuna, the major urban centers of the region, as well as heavy metals from fungicides used on cacao plantations (Klumpp *et al.* 2002; Lima *et al.* 2010).

Sampling Methods

The material examined was obtained mainly from February 2003 to February 2005 and in 2011 and 2012. The collection methods used from 2003 through 2005 were described by Almeida *et al.* (2006). In 2011 and 2012 the sampling was qualitative, with no standardization effort or predetermined frequency, and covered intertidal and shallow subtidal zones. Field activities were concentrated in the basins of the Cachoeira, Santana and Acuípe rivers. The caridean shrimps were sought in estuarine microhabitats such as burrows in fine sand and mud, on decomposing leaves and branches, on the roots and trunks of mangrove trees, under rocks or other hard substrates including oyster shells, and in association with algae and other invertebrates. The epibenthic shrimp were collected by hand or small hand nets in the intertidal or with hand nets in the shallow subtidal zone. We used a suction pump made of PVC pipe 50 mm in diameter to capture infaunal

species from mud and fine-sand bottoms. The sediment obtained with the pump was sieved to reveal the small-sized infaunal shrimp. Salinity, measured using a portable refractometer, ranged from 1 to 32.5 in this study. However, it may reach at least 36.5 at the mouth of the Cachoeira River (A.O. Almeida, pers. obs.). The collections conducted in this study complied with current applicable state and federal laws of Brazil (permanent license for collection of Zoological Material No. 24408-1 MMA/IBAMA/SISBIO for AOA).

The specimens were anesthetized on ice, fixed in 70% ethanol, and all but the specimens of *Alpheus brasiliensis* Anker, 2012 (specimen not collected, identified by means of the color pattern according to Anker 2012, p. 79, fig. 55, and p. 80, fig. 56) were deposited in the collection of crustaceans of the Universidade Estadual de Santa Cruz (UESC), Ilhéus. The classification adopted follows the proposal by De Grave and Fransen (2011). For complete synonymies of the taxa, also refer to De Grave and Fransen (2011). The order of species within families is alphabetical. The key for identification was developed taking as a starting point an adaptation of the keys proposed by Chace (1972) and Holthuis (1993). Additional distinctive characters were obtained by examining the material and in the original description of the taxa. Abbreviations used: (f) female, (m) male, (nov) non-ovigerous specimen, (ovf) ovigerous female.

RESULTS

Infraorder Caridea Dana, 1852

Superfamily Palaemonoidea Rafinesque, 1815

Family Palaemonidae Rafinesque, 1815

Leander paulensis Ortmann, 1897

Material examined: See material reported by Almeida et al. (2006).

Distribution: Western Atlantic – Florida, West Indies, and Brazil (Maranhão to São Paulo) (Ramos-Porto 1986). Previous records from Ilhéus: estuaries: Almeida et al. (2006; 2012); marine environment: Almeida et al. (2007a).

Macrobrachium acanthurus (Wiegmann, 1836)

Material examined: 1 m, 10 f (1 ovf), Santana River near Povoado do Rio de Engenho (14°51'09.3"S, 39°03'50.5"W), coll. A.O. Almeida, G.O. Soledade and P.S. Santos, 19.IV.2011 (UESC 1547); 5 m, Cachoeira River, Banco da Vitória, coll. F. Flores-Lopes, 2011 (UESC 1553). See also material reported by Almeida et al. (2006).

Distribution: North Carolina to Texas, Mexico, Cuba, Haiti, Dominican Republic, Puerto Rico, Nicaragua, Panama, Colombia, Venezuela, Suriname and Brazil (Pará to Rio Grande do Sul) (Melo 2003).

Previous records from Ilhéus: estuaries: Almeida et al. (2006, 2012); freshwater: Almeida et al. (2008) and Ferreira et al. (2010).

Palaemon northropi (Rankin, 1898)

Material examined: 6 m, 8 f, Cachoeira River, Banco da Sapetinga, confluence of Fundão and Cachoeira rivers (14°48'36.0"S, 39°02'38.9"W), coll. A.O. Almeida, G.O. Soledade and P.S. Santos, 19.IV.2011, salinity 2 (UESC




FIGURE 1. Main estuaries at Ilhéus, state of Bahia, Brazil. Gray area corresponds to Ilhéus urban area. (A) Sargi River, (B) Almada River, (C) Cachoeira River, (D) Santana River, (E) Cururupe River, (F) Acuípe River. Scale bar: 5 km.

1540); 1 f, Acuípe River (15°04'59.9"S, 38°59'55.9"W), coll. A.O. Almeida, G.O. Soledade and P.S. Santos, 23.VII.2009, on leaves on fine sand (UESC 1557). See also material reported by Almeida *et al.* (2006).

Distribution: Western Atlantic – Bermuda, West Indies, Central America, northern South America, Brazil (Ceará to Santa Catarina) and Uruguay (Ramos-Porto and Coelho 1990).

Previous records from Ilhéus: estuaries: Almeida *et al.* (2006) [as *Palaemon (Palaeander) northropi*]; marine environment: Almeida *et al.* (2012) (as *P. northropi*).

Palaemon pandaliformis (Stimpson, 1871)

Material examined: 1 f, Santana River, Fazenda Leoa (14°50'36.0"S, 39°02'46.7"W), coll. A.O. Almeida, G.O. Soledade and P.S. Santos, 19.IV.2011, salinity 1 (UESC 1546); 1 m, 6 f, Acuípe River (15°04'59.9"S, 38°59'55.9"W), coll. A.O. Almeida, G.O. Soledade and P.S. Santos, 23.VII.2009, on leaves on sand-mud bottom (UESC 1555). See also material reported by Almeida *et al.* (2006).

Distribution: Cuba, Puerto Rico, Barbados, Trinidad and Tobago, Guatemala, Nicaragua, Panama, Venezuela and Brazil (Rio Grande do Norte, and from Sergipe to Rio Grande do Sul, including Minas Gerais) (Melo 2003).

Previous records from Ilhéus: estuaries: Almeida *et al.* [2006, as *Palaemon (Palaemon) pandaliformis*; 2012, as *P. pandaliformis*]; freshwater: Almeida *et al.* (2008) [as *Palaemon (Palaemon) pandaliformis*].

Superfamily Alpheoidea Rafinesque, 1815

Family Alpheidae Rafinesque, 1815

Alpheus sp.

Material examined: See material reported by Almeida *et al.* (2006, as *A. heterochaelis* Say, 1818).

Previous records from Ilhéus: estuaries: Almeida *et al.* (2006, 2012, as *A. heterochaelis*).

Remarks: The material reported from Ilhéus in previous studies by Almeida *et al.* (2006, 2012) as *A. heterochaelis* is now recognized as a new species of *Alpheus* Fabricius, 1798, currently in process of description by the first author.

Alpheus angulosus McClure, 2002

Material examined: 7 m, 7 f, Cachoeira River, Maramata Beach (14°48'28.7"S, 39°01'33.3"W), coll. A.O. Almeida, A.M. Cunha, G.O. Soledade, M.V. Oliveira and P.S. Santos, 15.VII.2011, salinity 32.5, under rocks (UESC 1564); 1 m, Cachoeira River, Maramata Beach (14°48'28.7"S, 39°01'33.3"W), 23.III.2011, under rocks (UESC 1525).

Distribution: Western Atlantic - North Carolina to Florida, Gulf of Mexico, Caribbean Sea, French Guiana, and Brazil (Atol das Rocas, Fernando de Noronha, Ceará, Paraíba, Bahia, Rio de Janeiro, São Paulo, Santa Catarina, and Rio Grande do Sul) (Anker 2012).

Previous records from Ilhéus: estuaries: Almeida *et al.* (2013).

Alpheus bouvieri A. Milne-Edwards, 1878

Material examined: 8 m, 12 f, Cachoeira River, Maramata Beach (14°48'28.7"S, 39°01'33.3"W), coll. A.O. Almeida, A.M. Cunha, G.O. Soledade, M.V. Oliveira and P.S. Santos, 23.III.2011, salinity 7.5 (UESC 1558). See also material

reported by Almeida *et al.* (2006).

Distribution: Western Atlantic – Bermuda, Florida, West Indies, Panama, and Brazil (Atol das Rocas, Fernando de Noronha, and Ceará to Rio Grande do Sul). Central Atlantic - Ascension Island. Eastern Atlantic - Cape Verde, Senegal to Gulf of Guinea and Congo (Crosnier and Forest 1966; Christoffersen 1979, 1998; Manning and Chace 1990; Anker *et al.* 2009).

Previous records from Ilhéus: estuaries: Almeida *et al.* (2006; 2012).

Alpheus brasileiro Anker, 2012

Material examined: 1 ovf, not deposited, Cachoeira River, Av. Lomanto Jr. (14°48'31.1"S, 39°02'08.3"W), under rocks, 2012, recognized by color pattern (see Material and Methods section).

Distribution: Western Atlantic - Brazil (Pará, Ceará, Rio Grande do Norte, Alagoas, Bahia, Rio de Janeiro, São Paulo, and Santa Catarina) (Anker 2012).

Previous records from Ilhéus: none.

Alpheus buckupi Almeida, Terossi, Araújo-Silva and Mantelatto, 2013

Material examined: 2 m, Cachoeira River, Maramata Beach (14°48'28.7"S, 39°01'33.3"W), coll. A.O. Almeida, A.M. Cunha, G.O. Soledade, M.V. Oliveira and P.S. Santos, 23.III.2011, salinity 7.5, under rocks (UESC 1559).

Distribution: Western Atlantic – Venezuela (Orinoco Delta) and Brazil (Ceará, Rio Grande do Norte, Pernambuco, Alagoas, Bahia and São Paulo). Eastern Atlantic – São Tomé and Príncipe (Almeida *et al.* 2013).

Previous records from Ilhéus: none.

Alpheus carlae Anker, 2012

Material examined: 2 m, 4 f, mouth of Almada River, Station 2 (14°46'27.2"S, 39°03'14.8"W), coll. A.O. Almeida and J.T.A. Santos, 22.II.2005 (UESC 551); 1 m, 3 f, Cachoeira River, Station 8, Pontal, trawl (14°48'10"S, 39°02'12.3"W), coll. A.O. Almeida, J.T.A. Santos and J.R. Luz, 22.II.2005 (UESC 598); 1 f, Cachoeira River, Av. Lomanto Jr. (14°48'31.1"S, 39°02'08.3"W), coll. A.O. Almeida, G.O. Soledade, P.S. Santos and A.C. Costa-Souza, 18.IV.2011, salinity 14, burrow in mud (UESC 1551); 9 m, 2 f, Cachoeira River, Maramata Beach (14°48'28.7"S, 39°01'33.3"W), coll. A.O. Almeida, A.M. Cunha, G.O. Soledade, M.V. Oliveira and P.S. Santos, 17.V.2011, salinity 29, under rocks (UESC 1562); 1 ovf, Cachoeira River, Maramata Beach (14°48'28.7"S, 39°01'33.3"W), 04.II.2011, under rocks (UESC 1528).

Distribution: Western Atlantic - southern Florida, Puerto Rico, Jamaica, Belize, Panama, Venezuela, French Guiana, and Brazil (Ceará to São Paulo) (Anker 2012).

Previous records from Ilhéus: estuaries: Almeida *et al.* [2006, as *A. armillatus* (H. Milne Edwards, 1837 - in part; 2012, as *A. cf. armillatus* - in part; 2013, as *A. carlae*].

Alpheus chacei Carvacho, 1979

Material examined: 3 m, 2 f, Cachoeira River, Av. Lomanto Jr. (14°48'31.1"S, 39°02'08.3"W), coll. A.O. Almeida, G.O. Soledade and P.S. Santos, 18.IV.2011, salinity 14, burrows in mud (UESC 1548).

Distribution: Western Atlantic - French Antilles

(Guadeloupe) and Brazil (Pará and Paraíba to São Paulo) (Christoffersen 1979, as *A. maxilliplanus* Christoffersen, 1979; 1984; 1998; Soledade and Almeida 2013).

Previous records from Ilhéus: none.

Alpheus estuariensis Christoffersen, 1984

Material examined: 3 m, 2 f, Cachoeira River, Banco da Sapetinga, confluence of Cachoeira and Fundão rivers (14°48'36.0"S, 39°02'38.9"W), coll. A.O. Almeida, G.O. Soledade and P.S. Santos, 19.IV.2011, salinity 2, burrows in mud (UESC 1539); 2 f, Santana River, Nossa Senhora da Maré Island (14°50'57.8"S, 39°03'15.0"W), coll. A.O. Almeida, G.O. Soledade and P.S. Santos, 19.IV.2011, salinity 2 (UESC 1542); 2 m, 3 f, Santana River, Fazenda Leoa (14°50'36.0"S, 39°02'46.7"W), coll. A.O. Almeida, G.O. Soledade and P.S. Santos, 19.IV.2011, salinity 1, under rocks (UESC 1544); 1 f, Cachoeira River, Av. Lomanto Jr. (14°48'31.1"S, 39°02'08.3"W), coll. A.O. Almeida, G.O. Soledade, P.S. Santos and A.C. Costa-Souza, 18.IV.2011, salinity 14, burrows in mud (UESC 1550); 2 m, 2 f, Acuípe River (15°04'59.9"S, 38°59'55.9"W), coll. A.O. Almeida, G.O. Soledade and P.S. Santos, 23.VII.2009, under rocks (UESC 1556). See also material reported by Almeida *et al.* (2006).

Distribution: Western Atlantic – Florida, Mississippi to Texas, Cuba, Dominican Republic, Trinidad and Tobago, Curaçao, and Brazil (Pará to Santa Catarina) (Christoffersen 1984; Soledade and Almeida 2013; Almeida and Mantelatto in press).

Previous records from Ilhéus: estuaries: Almeida *et al.* (2006; 2012).

Alpheus intrinsecus Spence Bate, 1888

Material examined: See material reported by Almeida *et al.* (2006).

Distribution: Western Atlantic - Puerto Rico to Brazil (Piauí to Santa Catarina). Eastern Atlantic - Western Sahara to Gabon (Crosnier and Forest 1966; Christoffersen 1979; Soledade and Almeida 2013).

Previous records from Ilhéus: estuaries: Almeida *et al.* (2006, 2012); marine environment: Almeida *et al.* (2007a).

Alpheus nuttingi (Schmitt, 1924)

Material examined: 1 m, 1 f, Cachoeira River, Maramata Beach (14°48'28.7"S, 39°01'33.3"W), coll. A.O. Almeida, A.M. Cunha, G.O. Soledade, M.V. Oliveira and P.S. Santos, 20.IV.2011, salinity 10.5, under rocks (UESC 1561).

Distribution: Western Atlantic - southern Florida, southwestern Gulf of Mexico, and West Indies to Brazil (Ceará to Santa Catarina) (Anker *et al.* 2007; Soledade and Almeida 2013).

Previous records from Ilhéus: none.

Alpheus pontederiae de Rochebrune, 1883

Material examined: 1 f, Santana River, Nossa Senhora da Maré Island (14°50'57.8"S, 39°03'15.0"W), coll. A.O. Almeida, G.O. Soledade and P.S. Santos, 19.IV.2011, salinity 2 (UESC 1543); 1 m, 4 f, Santana River, Fazenda Leoa (14°50'36.0"S, 39°02'46.7"W), coll. A.O. Almeida, G.O. Soledade and P.S. Santos, 19.IV.2011, salinity 1, under rocks (UESC 1545). See also material reported by Almeida *et al.* (2006).

Distribution: Western Atlantic - Venezuela to Brazil (Pará, Maranhão, Paraíba, Alagoas, Bahia, São Paulo, Paraná). Eastern Atlantic - Senegal to Congo (Crosnier and Forest 1966; Christoffersen 1984; Soledade and Almeida 2013).

Previous records from Ilhéus: estuaries: Almeida *et al.* (2006, 2012).

Automate cf. dolichognatha De Man, 1888

(Figure 2)

Material examined: 1 nov, Cachoeira River, Maramata Beach (14°48'28.7"S, 39°01'33.3"W), coll. A.O. Almeida, A.M. Cunha, G.O. Soledade, M.V. Oliveira and P.S. Santos, 25.IX.2011, salinity 35, under rocks (UESC 1563). Additional material: 5 nov, Bahia, Itacaré, Concha Beach, mouth of Contas River (14°16'31.4"S, 38°59'14.5"W), coll. G.O. Soledade, P.S. Santos and M.V. Oliveira, 27.XI.2012, under rocks (UESC 1565).

Distribution: Western Atlantic – North Carolina, Yucatan Peninsula, Virgin Islands, Antigua, Barbados, Brazil (Rio de Janeiro). Central Atlantic – Ascension. Indo-West Pacific – Red Sea to Samoa. Eastern Pacific – Tres Marias, Revillagigedo, Cocos and Galapagos Islands, California, Gulf of California to Peru (Chace 1972, as *A. gardineri* Coutière, 1902; Banner and Banner 1973; Manning and Chace 1990; Christoffersen 1998; Wicksten and Hendrickx 2003).

Previous records from Bahia: none.

Remarks: *Automate dolichognatha*, originally described from Indonesia, is a widely distributed, almost pantropical species complex (Anker 2001; Anker and Komai 2004) and is in need of a comprehensive revision. The record of the species from Rio de Janeiro by Christoffersen (1998), in a list of alpheid shrimps from Brazil, is the only known report from the Brazilian coast. No illustrations or morphological account of this material were provided by Christoffersen (1998). Anker and Komai (2004) divided the genus *Automate* De Man, 1888 into three informal groups of species: *A. dolichognatha*, *A. evermanni* Rathbun, 1901, and *A. hayashii* Anker and Komai, 2004. The combination of characteristics such as (1) major chela subrectangular (see Figure 2D), (2) propodus of pereopod 3 with row of spiniform setae, (3) dactylus of pereopod 3-5 simple, subconical and (4) diaeresis of uropodal exopodite with two dorsal teeth place our specimens in the *A. dolichognatha* group. The other species within this group is the little-known eastern-Atlantic *A. talismani* Coutière, 1902, whose taxonomic status remains unclear (see Chace 1988; Anker and Komai 2004). For this reason our material is referred provisionally as *A. cf. dolichognatha*. Our material also agrees, in general, with the illustrations provided by Chace (1972) as *A. gardineri*. The color pattern of the Brazilian material is illustrated for the first time (Figure 2). The record of color pattern of fresh material will aid in mapping the geographic distribution of the taxa, after a taxonomic revision of *A. dolichognatha sensu lato* can be conducted and our material properly attributed.

Leptalpheus axianassae Dworschak and Coelho, 1999

Material examined: 1 m, 1 f, Cachoeira River, Av. Lomanto Jr. (14°48'31.1"S, 39°02'08.3"W), coll. A.O. Almeida, G.O. Soledade and P.S. Santos, 18.IV.2011, salinity




FIGURE 2. *Automate cf. dolichognatha* De Man, 1888, non-ovigerous specimens, Itacaré, Bahia, Brazil (UESC 1565). A, habitus, dorsal view; B, habitus, lateral view; C, anterior region, lateral view; D, major chela (above), minor chela (below), lateral view. Scale bars: 1 mm.

14, burrows in mud (UESC 1552).

Distribution: Western Atlantic – Florida and Brazil (Pernambuco, Bahia and São Paulo) (Dworschak and Coelho 1999; Anker *et al.* 2006a; Almeida *et al.* 2012).

Previous records from Ilhéus: none.

Salmones carvachoi Anker, 2007

Material examined: 1 nov, Cachoeira River, Banco da Sapetinga, confluence of Cachoeira and Fundão rivers (14°48'36.0"S, 39°02'38.9"W), coll. A.O. Almeida, G.O. Soledade and P.S. Santos, 19.IV.2011, salinity 2, burrows in mud (UESC 1541); 4 ovf, Cachoeira River, Av. Lomanto Jr. (14°48'31.1"S, 39°02'08.3"W), coll. A.O. Almeida, G.O. Soledade, P.S. Santos and A.C. Costa-Souza, 18.IV.2011, salinity 14, burrows in mud (UESC 1549).

Distribution: Western Atlantic – Mexico (Yucatan), Guadeloupe and Brazil (Paraíba, Pernambuco, Bahia, São Paulo and Paraná) (Anker 2007; 2010).

Previous records from Ilhéus: none.

Synalpheus fritzmuelleri Coutière, 1909

Material examined: See material reported by Almeida *et al.* (2006).

Distribution: Western Atlantic – Bermuda, North Carolina to Florida, Gulf of Mexico, Bahamas, Caribbean Sea (Yucatan to Venezuela) and Brazil (São Pedro and São Paulo Archipelago, Atol das Rocas, Ceará to Santa Catarina). Central Atlantic – Ascension and Saint Helena islands (Christoffersen 1979, 1998; Holthuis *et al.* 1980;

Manning and Chace 1990; Anker *et al.* 2012).

Previous records from Ilhéus: estuaries: Almeida *et al.* (2006, 2012).

Family Hippolytidae Spence Bate, 1888

Latreutes parvulus (Stimpson, 1871)

Material examined: 1 f, Ilhéus, Cachoeira River, Station 8, Pontal, trawl (14°48'10"S, 39°02'12.3"W), coll. A.O. Almeida, J.T.A. Santos, N.R. Ferraz and C.S. Soares, 18.IX.2004 (UESC 589).

Distribution: Western Atlantic – New Jersey, USA, to Buenos Aires, Argentina. Eastern Atlantic – Western Sahara to Congo (Christoffersen 1982).

Previous records from Ilhéus: estuaries: Almeida *et al.* (2012).

Merguia rhizophorae (Rathbun, 1900)

Material examined: 4 nov, Cachoeira River, Maramata Beach (14°48'28.7"S, 39°01'33.3"W), coll. A.O. Almeida, A.M. Cunha, G.O. Soledade, M.V. Oliveira and P.S. Santos, 23.III.2011, salinity 7.5 (UESC 1560). See also material reported by Almeida *et al.* (2006).

Distribution: Western Atlantic – Panama, Suriname and Brazil (Piauí to Bahia). Eastern Atlantic – Niger Delta in Nigeria (Chace 1972; Bruce 1993; Christoffersen 1998; Almeida *et al.* 2006).

Previous records from Ilhéus: estuaries: Almeida *et al.* (2006, 2012).

Family Ogyrididae Holthuis, 1955
Ogyrides alphaerostris (Kingsley, 1880)
Material examined: 1 f, mouth of Acuípe River (15°04'59.9"S, 38°59'55.9"W), coll. A.O. Almeida, G.O. Soledade and P.S. Santos, 23.VII.2009, on mud (UESC 1554).
Distribution: Western Atlantic – Virginia to Brazil (Amapá, Pará, Paraíba to Alagoas, Rio de Janeiro to Rio Grande do Sul) (Williams 1984; Christoffersen 1998; Coelho *et al.* 2006).
Previous records from Bahia: none.

Key for identification of estuarine caridean shrimps from Ilhéus:

- 1. Carpus of the second pereopod not subdivided (Palaemonidae) 2
- 1'. Carpus of the second pereopod subdivided 5

- 2. Carapace with hepatic spine. *Macrobrachium acanthurus*
- 2'. Carapace without hepatic spine 3

- 3. Branchiostegal groove between antennal and branchiostegal spines absent; endopod of male first pleopod with well-developed appendix interna; propodus of fifth pereopod without transverse rows of grooming setae on distal part of posterior margin. .. *Leander paulensis*
- 3'. Branchiostegal groove between antennal and branchiostegal spines present; endopod of male first pleopod without appendix interna; propodus of fifth pereopod with transverse rows of grooming setae on distal part of posterior margin 4

- 4. Carpus of second pereopod as long as chela; rostrum with 3-4 teeth on ventral margin *Palaemon northropi*
- 4'. Carpus of second pereopod twice as long as chela; rostrum with 5-8 teeth (rarely 9) on ventral margin
..... *Palaemon pandaliformis*

- 5. Eyestalks very long, reaching nearly to distal margin of antennular peduncle; first pereopods about as robust as second pereopods (Ogyrididae) *Ogyrides alphaerostris*
- 5'. Eyestalks not unusually long, not reaching beyond distal margin of first segment of antennular peduncle, sometimes covered by carapace; first pereopods more robust than second pereopods 6

- 6. Carapace without cardiac notch on posterior margin; first pereopods usually equal, not swollen; eyes totally exposed and freely movable; rostrum well-developed, toothed (Hippolytidae) 7
- 6'. Carapace with cardiac notch on posterior margin; first pereopods often unequal and swollen; eyes often totally or partially covered by ocular hoods; rostrum relatively short, smooth-edged (Alpheidae) 8

- 7. Merus of second pereopod not subdivided, carpus subdivided into three segments *Latreutes parvulus*
- 7'. Merus and carpus of second pereopod multiarticulate (24-27 and 10-14 segments, respectively)
..... *Merguia rhizophorae*

- 8. Posterolateral angle of sixth abdominal segment with movable triangular plate *Leptalpheus axianassae*
- 8'. Posterolateral angle of sixth abdominal segment without movable triangular plate 9

- 9. Eyes not concealed in anterior view; dactylus of major chela without a large molar tooth (plunger) fitting in deep fossa on pollex 10
- 9'. Eyes totally covered by ocular hoods from all but anteroventral view; dactylus of major chela with a large molar tooth (plunger) fitting in deep fossa on pollex 11

- 10. Eyes, including peduncle, totally exposed dorsally; rostral projection not reaching beyond anterolateral margin of carapace *Automate cf. dolichognatha*
- 10'. Eyes covered in dorsal view; distal tip of rostrum reaching beyond anterolateral margin of carapace
..... *Salmoneus carvachoi*

- 11. Pereiopods without epipods; second pleopod of male without appendix masculina; ocular hoods with tooth on anterior margin (front tridentate)
..... *Synalpheus fritzmulleri*
- 11'. Pereiopods 1-4 with epipods; second pleopod of male without appendix masculina; ocular hoods without tooth on anterior margin (rounded) 12

- 12. Rostrum dorsally flat; ocular hoods with tooth arising from surface of mesial slope, overhanging adrostral furrows *Alpheus intrinsecus*
- 12'. Rostrum rounded or carinate in dorsal midline, not flattened; ocular hoods unarmed 13

- 13. Merus of chelipeds unarmed at distal end of ventromesial margin 14
- 13'. Merus of first chelipeds armed with sharp tooth at distal end of ventromesial margin 19

- 14. Dactylus of third and fourth pereiopods conical 15
- 14'. Dactylus of third and fourth pereiopods usually subspatulate 17

- 15. Dorsal groove of major chela palm not extending posteriorly on mesial surface *Alpheus bouvieri*
- 15'. Dorsal groove of major chela palm extending posteriorly on mesial surface 16

- 16. Rostral carina slightly broadening posteriorly; ventromesial carina of first segment with acute tooth, directed forward (hook-shaped); fingers of minor chela without balaeniceps setae in male; female minor chela more robust (propodus approx. 2.7 times as long as broad) *Alpheus nuttingi*
- 16'. Rostral carina not broadening posteriorly; ventromesial carina of first segment with rounded tooth; fingers of minor chela with balaeniceps setae in male; female minor chela more slender (propodus approx. 4 times as long as broad) *Alpheus* sp.

- 17. Distal margin of pollex of major chela with rounded ending; fingers of minor chela with balaeniceps setae

in males; uropodal exopod with two sharp teeth on posterolateral margin, one on each side of spiniform setae	<i>Alpheus pontederiae</i>
17'. Distal margin of pollex of major chela distinctly truncate; fingers of minor chela without balaeniceps setae in both males and females; uropodal exopod with one sharp tooth on posterolateral margin, lateral to spiniform setae	18
18. Fingers of minor chela distinctly longer than palm; antepenultimate segment of third maxilliped broadly enlarged	<i>Alpheus chacei</i>
18'. Fingers of minor chela as long as palm; antepenultimate segment of third maxilliped not enlarged	<i>Alpheus estuariensis</i>
19. Adrostral furrows shallow, not sharply delimited; fingers of minor chela of male with balaeniceps setae	<i>Alpheus buckupi</i>
19'. Adrostral furrows deep, sharply delimited; fingers of minor chela of male without balaeniceps setae	20
20. Major chela with dactylus plunger relatively short, its proximal height at most 0.6 length of distolateral margin; dactylus plunger orange-colored mesially (fresh specimens)	<i>Alpheus brasileiro</i>
20'. Major chela with dactylus plunger relatively long, its proximal height more than 0.7 length of distolateral margin; dactyls plunger whitish mesially (fresh specimens)	21
21. First and second abdominal sternite with strong median process in males; fifth pereopod ischium with or without spiniform seta on ventrolateral surface	<i>Alpheus carlae</i>
21'. First abdominal sternite with small median process, second abdominal sternite unarmed in males; fifth pereopod ischium always unarmed on ventrolateral surface	<i>Alpheus angulosus</i>

DISCUSSION

The list of caridean shrimps from estuaries of Ilhéus currently comprises 22 species in four families. It is difficult to compare species richness among estuaries because the number of species recorded in surveys may vary depending on several factors such as the size of the study area, climate and oceanographic conditions, sampling effort, diversity of substrata available, type(s) of substrata sampled, method(s) of sampling, and the experience and taxonomic group of expertise of the collector(s), among others.

The number of caridean species obtained in this study is high compared to the numbers obtained in some quantitative studies carried out in estuaries or

other shallow-water coastal areas in both warm and cold temperate areas (e.g., Gore et al. 1981; Able et al. 2002; Neves et al. 2007) (see Table 1). This result was expectable, since there is a general trend toward a decrease in the number of caridean species at higher latitudes (d’Udekem d’Acoz 1999; Boschi 2002; Clarke and Crame 2010), as observed for other marine groups. With a total of 14 species, the alpheidids were the most prominent family with respect to species richness. Indeed, alpheid shrimps are highly diverse in tropical regions, where they occupy a wide variety of marine and estuarine habitats, living on various types of bottom and in association with other phyla of invertebrates as well as with fish (Anker et al. 2006b). In contrast, the richness of palaemonids and hippolytids is proportionally higher in shallow-water environments in temperate areas, especially in areas covered with vegetation (Hooks et al. 1976; Gore et al. 1981; López de la Rosa et al. 2002; Glancy et al. 2003).

The number of species recorded in estuaries of Ilhéus is lower than that reported by Carvacho (1979) (n=29) in mangroves from Guadeloupe (Antillean Province, see Boschi 2000) and higher than the numbers reported by Hendrickx (1984) (n=4) and Echeverría-Sáenz et al. (2003) (n=11) in estuaries of the Mexican and Costa Rican Pacific coasts (Panamic Province, see Boschi 2000) (see Table 1). The number of species reported by Carvacho (1979) would be expected to outnumber those in the present study, because the Antillean Province is the richest in number of decapod species among all the zoogeographic marine provinces of the Americas (see Boschi 2000).

The estuarine caridean fauna along the northern and northeastern Brazilian coast has been very little studied. The epibenthic carideans (e.g., most palaemonids and hippolytids) are in general well documented. On the other hand, the (ecologically) cryptic fauna is especially little known, especially the soft-bottom infauna. The difficulties of collection and accurate taxonomic identification partly explain this situation, which is not exclusive to this area. The number of species recorded here is much higher than the numbers found in the vast majority of surveys that have been carried out along the Brazilian coast (e.g., Ramos-Porto 1980, Ilha de Itamaracá, Pernambuco; Coelho-Santos and Coelho 2001, Paripe River, Pernambuco; Calado and Sousa 2003, Alagoas; Ferreira and Sankarankutty 2003, Rio Grande do Norte), most of them general studies on decapod fauna. The sampling effort concentrated on carideans, allied to recent advances in the taxonomy of certain taxa (e.g. Anker 2012; Almeida et al. 2013) are the main factors responsible for this relatively long species list. Systematic surveys in other estuaries along this stretch of the Brazilian coast are likely to reveal local caridean faunas as rich as or richer than that recorded from Ilhéus. Sampling of the infauna is especially likely to result in the discovery of new records and new taxa for science.



TABLE 1. Comparison of the composition of the caridean fauna in several estuarine regions.

STUDY AREA	NUMBER OF SPECIES	REFERENCE
Terraba-Sierpe mangrove system, Costa Rica, Pacific side	11 (1 atyid, 6 palaemonids, 4 alpheids)	Echeverría-Sáez et al. (2003)
Estero El Verde, Sinaloa, Mexico, Pacific side	3 (2 palaemonids, 1 alpheid)	Hendrickx (1984)
Cape Cod estuary, Massachusetts, USA	3 (1 palaemonid, 1 crangonid, 1 hippolytid)	Able et al. (2002)
Indian River Lagoon estuary, Florida, USA	13 (3 palaemonids, 4 alpheids, 4 hippolytids, 2 processids)	Gore et al. (1981)
off mouths of Encofina and Fenholloway rivers, Apalachee Bay, Florida, USA	19 (7 palaemonids, 5 alpheids, 6 hippolytids, 1 processid)	Hooks et al. (1976)
Mangroves from Guadeloupe, the Caribbean	29 (1 atyid, 9 palaemonids, 11 alpheids, 1 ogyridid, 5 hippolytids, 2 processids)	Carvacho (1979)
Estuaries of Casqueira, Conceição and Potengi rivers, Rio Grande do Norte, Brazil	12 (4 palaemonids, 4 alpheids, 1 hippolytid, 1 ogyridid, 1 processid)	Ferreira and Sankarankutty (2002)
Ilha de Itamaracá, Pernambuco, Brazil	14 (1 atyid, 6 palaemonids, 3 alpheids, 4 hippolytids) (only estuarine species)	Ramos-Porto (1980)
Paripe River estuary, Pernambuco, Brazil	11 (1 atyid, 6 palaemonids, 3 alpheids, 1 hippolytid)	Coelho-Santos and Coelho (2001)
Mundau/Manguaba estuary-lagoon system, Alagoas, Brazil	13 (1 atyid, 6 palaemonids, 6 alpheids)	Calado and Sousa (2003)
Camamu Bay, Bahia, Brazil	22 (1 pasiphaeid, 5 palaemonids, 11 alpheids, 5 hippolytids)	Almeida et al. (2007b)
Estuaries of Ilhéus, Bahia, Brazil	22 (4 palaemonids, 15 alpheids, 2 hippolytids, 1 ogyridid)	Present study
Sado estuary, Portugal	6 (2 palaemonids, 1 alpheid, 1 hippolytid, 1 processid, 1 crangonid)	Neves et al. (2007)
Valdelagrana Beach, off Guadalete River mouth, Bay of Cadiz, Spain	20 (3 palaemonids, 1 alpheid, 6 hippolytids, 1 ogyridid, 5 processids, 4 crangonids)	López de la Rosa et al. (2002)

ACKNOWLEDGMENTS: The authors are indebted to the Fundação de Amparo à Pesquisa do Estado da Bahia (FAPESB) (PPP0073/2010; APP 0035/2011) and to the Universidade Estadual de Santa Cruz (00220.1100.590; 00220.1100.1065) for supporting the projects “Diversidade de Crustáceos Decápodos Marinhos e Estuarinos do Sul da Bahia, Brasil” and “Biologia e Ecologia de Camarões da Família Alpheidae (Crustacea: Decapoda: Caridea)”. To Janet W. Reid for assistance with the English text. To two anonymous referees for the criticisms that greatly improved the manuscript. ACCS, PSS and GOS thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for the provision of a Master’s scholarship. AMC thanks the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for the provision of a scientific initiation scholarship.

LITERATURE CITED

Able, K.W., M.P. Fahay, K.L. Heck Jr., C.T. Roman, M.A. Lazzari and S.C. Kaiser, 2002. Seasonal distribution and abundance of fishes and decapod crustaceans in a Cape Cod estuary. *Northeastern Naturalist* 9(3): 285-302.

Almeida, A.O., G. Boehs, C.L. Araújo-Silva and L.E.A. Bezerra. 2012. Shallow-water caridean shrimps from southern Bahia, Brazil, including the first record of *Synalpheus ul* (Ríos & Duffy, 2007) (Alpheidae) in the southwestern Atlantic Ocean. *Zootaxa* 3347: 1-35.

Almeida, A.O., P.A. Coelho, J.R. Luz, J.T.A. Santos and N.R. Ferraz. 2008. Decapod crustaceans in fresh waters of southeastern Bahia, Brazil. *Revista de Biología Tropical* 56(3): 1225-1254.

Almeida, A.O., P.A. Coelho, J.T.A. Santos and N.R. Ferraz. 2006. Crustáceos decápodos estuarinos de Ilhéus, Bahia, Brasil. *Biota Neotropica* 6(2): 1-24.

Almeida, A.O., P.A. Coelho, J.T.A. Santos and N.R. Ferraz. 2007a. Crustáceos estomatópodos e decápodos da costa de Ilhéus, Bahia, Brasil. *Atlântica* 29(1): 5-20.

Almeida, A.O., M.C. Guerrazzi and P.A. Coelho. 2007b. Stomatopod and decapod crustaceans from Camamu Bay, state of Bahia, Brazil. *Zootaxa* 1553: 1-45.

Almeida, A.O. and F.L. Mantelatto. In press. Extension of the known distributions of three estuarine snapping shrimps of the genus *Alpheus* Fabricius, 1798 (Caridea, Alpheidae) in South America. *Crustaceana*.

Almeida, A.O., M. Terossi, C.L. Araújo-Silva and F.L. Mantelatto. 2013. Description of *Alpheus buckupi* spec. nov., a new amphi-Atlantic snapping shrimp (Caridea: Alpheidae), based on morphological and molecular data. *Zootaxa* 3652: 437-452.

Andrade, M.P. 2003. *Ilhéus, passado e presente*. Ilhéus: Editus. 144p.

Anker, A. 2001. Two new species of snapping shrimps from the Indo-Pacific, with remarks on colour patterns and sibling species in Alpheidae (Crustacea: Caridea). *The Raffles Bulletin of Zoology* 49(1): 57-72.

Anker, A. 2007. New species and records of alpheid shrimps, genera *Salmonus* Holthuis and *Parabetaeus* Coutière, from the tropical western Atlantic (Decapoda, Caridea). *Zootaxa* 1653: 21-39.

Anker, A. 2010. The shrimp genus *Salmonus* Holthuis, 1955 (Crustacea, Decapoda, Alpheidae) in the tropical western Atlantic, with description of five new species. *Zootaxa* 2372: 177-205.

Anker, A. 2012. Revision of the western Atlantic members of the *Alpheus armillatus* H. Milne Edwards, 1837 species complex (Decapoda, Alpheidae), with description of seven new species. *Zootaxa* 3386: 1-109.

Anker, A., S.T. Ah Yong, P.Y. Noël and A.R. Palmer. 2006b. Morphological phylogeny of alpheid shrimps: parallel preadaptation and the origin of a key morphological innovation, the snapping claw. *Evolution* 60 (12): 2507-2528.

Anker, A., C. Hurt and N. Knowlton. 2007. Revision of the *Alpheus nuttingi* (Schmitt) species complex (Crustacea: Decapoda: Alpheidae), with description of a new species from the tropical eastern Pacific. *Zootaxa* 1577: 41-60.

Anker, A., C. Hurt and N. Knowlton. 2009. Description of cryptic taxa within the *Alpheus bouvieri* A. Milne-Edwards, 1878 and *A. hebes* Abele & Kim, 1988 species complexes (Crustacea: Decapoda: Alpheidae). *Zootaxa* 2153: 1-23.

Anker, A. and T. Komai. 2004. Descriptions of two new species of alpheid shrimps from Japan and Australia, with notes on taxonomy of *Automate* De Man, *Coronalpheus* Wicksten and *Bermudacaris* Anker and Iliffe (Crustacea: Decapoda: Caridea). *Journal of Natural History* 38(15): 1895-1915.

Anker, A., P.P.G. Pachelle, S. De Grave and K.M. Hultgren. 2012. Taxonomic and biological notes on some Atlantic species of the snapping shrimp genus *Synalpheus* Spence Bate, 1888 (Decapoda, Alpheidae). *Zootaxa* 3598: 1-96.

Anker, A., J. Vera Caripe and C. Lira. 2006a. Description of a new species of commensal alpheid shrimp (Crustacea, Decapoda) from the southern Caribbean Sea. *Zoosystema* 28(3): 683-702.

Banner, D.M. and A.H. Banner. 1973. The alpheid shrimp of Australia. Part I. The lower genera. *Records of the Australian Museum* 28(15): 291-382.

Bauer, R.T. 2004. *Remarkable shrimps: adaptations and natural history of the carideans*. Norman: University of Oklahoma Press. 282 p.

Boschi, E.E. 2000. Species of decapod crustaceans and their distribution in the American marine zoogeographic provinces. *Revista de Investigación y Desarrollo Pesquero* 13: 7-136.

Boschi, E.E. 2002. Distribution of Continental Shelf Decapod Crustaceans Along the American Pacific Coast; p. 235-239 In E. Escobar-Briones and F. Alvarez (eds). *Modern Approaches to the Study of Crustacea*. New York: Kluwer Academic / Plenum Publishers.

Bruce, A.G. 1993. The occurrence of the semi-terrestrial shrimps *Merguia oligodon* (De Man 1888) and *M. rhizophorae* (Rathbun 1900)

- (Crustacea Decapoda Hippolytidae) in Africa. *Tropical Zoology* 3: 179-187.
- Calado, T.C.S. and E.C. Sousa. 2003. *Crustáceos do complexo estuarino-lagunar Mundaú/Manguaba – Alagoas*. Maceió: FAPEAL. 116 p.
- Carvacho, A. 1979. Les crevettes carides de la mangrove guadeloupéenne. *Bulletin du Muséum national d'Histoire naturelle* (4), section A, Zoologie, Biologie et Écologie animales 1: 445-470.
- Chace Jr., F.A. 1972. The shrimps of the Smithsonian-Bredin Caribbean Expeditions with a summary of the West Indian shallow-water species (Crustacea: Decapoda: Natantia). *Smithsonian Contributions to Zoology* 98: 1-179.
- Chace Jr., F.A. 1988. The caridean shrimps (Crustacea: Decapoda) of the Albatross Philippine Expedition, 1907-1910, Part 5: Family Alpheidae. *Smithsonian Contributions to Zoology* 466: 1-99.
- Christoffersen, M.L. 1979. Decapod Crustacea: Alpheoidea. Campagne de la Calypso au large des côtes atlantiques de l'Amérique du Sud (1961-1962). I. 36, *Annales de l'Institut Océanographique, Monaco*, (Suppl.) 55: 297-377.
- Christoffersen, M.L. 1982. Geographic distribution of warm water alpheoid shrimp (Crustacea, Caridea) on the continental shelf of eastern South America between 23 and 35° Lat. S. *Boletim do Instituto Oceanográfico* 31(1): 93-112.
- Christoffersen, M.L. 1984. The western Atlantic snapping shrimps related to *Alpheus heterochaelis* Say (Crustacea, Caridea), with the description of a new species. *Papéis Avulsos de Zoologia* 35(19): 189-208.
- Christoffersen, M.L. 1998. Malacostraca. Eucarida. Caridea. Crangonoidea and Alpheoidea (Except Glyphocrangonidae and Crangonidae); p. 351-372 In P.S. Young (ed.). *Catalogue of Crustacea of Brazil*. Rio de Janeiro: Museu Nacional.
- Clarke, A. and J.A. Crame 2010. Evolutionary dynamics at high latitudes: speciation and extinction in polar marine faunas. *Philosophical Transactions of the Royal Society B*: 365: 3655-3666.
- Coelho, P.A., A.O. Almeida, J.F. Souza-Filho, L.E.A. Bezerra and B.W. Giraldez. 2006. Diversity and distribution of the marine and estuarine shrimps (Dendrobranchiata, Stenopodidea and Caridea) from North and Northeast Brazil. *Zootaxa* 1221: 41-62.
- Coelho-Santos, M.A. and P.A. Coelho. 2001. Crustacea Decapoda of the Paripe River Estuary, Pernambuco, Brazil. *Hydrobiologia* 449: 77-79.
- Coutière, H., 1902. Sur quelques espèces nouvelles du genre *Automate* De Man. *Bulletin du Muséum d'Histoire Naturelle* 8: 337-342.
- Coutière, H. 1909. The American species of the snapping shrimps of the genus *Synalpheus*. *Proceedings of the United States National Museum* 36: 1-93.
- Crosnier, A. and J. Forest. 1966. Crustacés Décapodes: Alpheidae. In Campagne de la Calypso dans le Golfe de Guinée et aux Iles Principe, São Tomé et Annobon (1956), et campagne aux Iles du Cap Vert (1959) (suite). Parte 19, Résultats Scientifiques des Campagnes de la Calypso, 7(27). *Annales de l'Institut Océanographique, Monaco* 44: 199-314.
- Dana, J.D. 1852. *United States Exploring Expedition during the years 1838, 1839, 1840, 1841, 1842, under the Command of Charles Wilkes, U.S.N.* Volume 13. Crustacea. Part I: 1-685, 1-27, Plates 1-96 (1855). Philadelphia: C. Sherman.
- De Grave, S., Y. Cai and A. Anker. 2008. Global diversity of shrimps (Crustacea: Decapoda: Caridea) in freshwater. *Hydrobiologia* 595: 287-293.
- De Grave, S. and C.H.J.M. Fransen. 2011. Carideorum catalogus: The Recent species of the dendrobranchiate, stenopodidean, procarididean and caridean shrimps (Crustacea: Decapoda). *Zoologische Mededelingen* 85(9): 195-589.
- Dworschak, P.C. and V.R. Coelho. 1999. On two alpheids from Araújo (São Paulo, Brazil) with a description of a new species of *Leptalpheus* (Decapoda: Caridea: Alpheidae). *Annalen des Naturhistorischen Museums in Wien* 101B: 475-488.
- Echeverría-Sáenz, S., R. Vargas and I.S. Wehrmann. 2003. Diversity of decapods inhabiting the largest mangrove system of Pacific Costa Rica. *Nauplius* 11(2): 91-97.
- Eggleston, D.B., L.L. Etherington and W.E. Elis. 1998. Organism response to habitat patchiness: species and habitat-dependent recruitment of decapod crustaceans. *Journal of Experimental Marine Biology and Ecology* 223: 111-132.
- Fabricius, J.C. 1798. *Supplementum Entomologiae Systematicae*, p. 1-572. Hafniae: Proft et Storch.
- Ferreira, A.C. and C. Sankarankutty. 2002. Estuarine carcinofauna (Decapoda) of Rio Grande do Norte, Brazil. *Nauplius* 10: 121-129.
- Ferreira, R.S., R.R.R. Vieira and F. D'Incao. 2010. The marine and estuarine shrimps of the Palaemoninae (Crustacea: Decapoda: Caridea) from Brazil. *Zootaxa* 2606: 1-24.
- Glancy, T.P., T.K. Frazer, C.E. Cichra and W.J. Lindberg. 2003. Comparative Patterns of Occupancy by Decapod Crustaceans in Seagrass, Oyster, and Marsh-edge Habitats in a Northeast Gulf of Mexico Estuary. *Estuaries* 26(5): 1291-1301.
- Gore, R.H., E.E. Gallaher, L.E. Scotto and W.A. Wilson. 1981. Studies of Decapod Crustacea from the Indian River Region of Florida. XI. Community Composition, Structure, Biomass and Species-Areal Relationships of Seagrass and Drift Algae-associated macrocrustaceans. *Estuarine, Coastal and Shelf Science* 12: 485-508.
- Hendrickx, M.E. 1984. Studies of the coastal marine fauna of southern Sinaloa, Mexico. II. The Decapod Crustaceans of Estero el Verde. *Anales del Instituto de Ciencias del Mar y Limnología, Universidad Autónoma, México* 11(1): 23-48.
- Holthuis, L.B. 1955. The recent genera of the caridean and stenopodidean shrimps (Class Crustacea, order Decapoda, supersection Natantia) with keys for their determination. *Zoologische Verhandelingen* 26: 1-157.
- Holthuis, L.B. 1993. *The recent genera of the caridean and stenopodidean shrimps (Crustacea, Decapoda with an appendix on the order Amphionidacea*. Leiden: Nationaal Natuurhistorisch Museum. 328 p.
- Holthuis, L.B., A.J. Edwards and H.R. Lubbock. 1980. The Decapod and Stomatopod Crustacea of St Paul's Rocks. *Zoologische Mededelingen* 56(3): 27-51.
- Hooks, T.A., K.L. Heck Jr. and R.L. Livingston. 1976. An inshore marine invertebrate community: structure and habitat associations in the northeastern Gulf of Mexico. *Bulletin of Marine Science* 26(1): 99-109.
- Kingsley, J.S. 1878. Notes on the North American Caridea in the Museum of the Peabody Academy of Science at Salem, Mass. *Proceedings of the Academy of Natural Sciences of Philadelphia* 30: 89-98.
- Kingsley, J.S. 1880. On a collection of Crustacea from Virginia, North Carolina, and Florida, with a revision of the genera of Crangonidae and Palaemonidae. *Proceedings of the Academy of Natural Sciences of Philadelphia* 31(3): 383-427, plate 14.
- Klumpp, A., K. Bauer, C. Franz-Gerstein and M. Menezes. 2002. Variation of nutrient and metal concentrations in aquatic macrophytes along the Rio Cachoeira in Bahia. *Environment International* 28: 165-171.
- Lima, M.C., M.F.L. Souza, G.F. Eça and M.A.M. Silva. 2010. Export and retention of dissolved inorganic nutrients in the Cachoeira River, Ilhéus, Bahia, Brazil. *Journal of Limnology* 69(1): 138-145.
- López de la Rosa, I., J.E. García Raso and A. Rodríguez. 2002. Evolution of a decapod community (Crustacea) of shallow soft bottoms with seaweeds from southern Europe. *Journal of the Marine Biological Association of the United Kingdom* 82: 85-95.
- Man, J.G. de. 1888. Bericht über die von Herrn Dr. J. Brock im indischen Archipel gesammelten Decapoden und Stomatopoden. *Archiv für Naturgeschichte* 53(3): 289-600, Plates 11-22a.
- Manning, R.B. and F.A. Chace Jr. 1990. Decapod and Stomatopod Crustacea from Ascension Island, South Atlantic Ocean. *Smithsonian Contributions to Zoology* 503: 1-91.
- McClure, M. 2002. Revised nomenclature of *Alpheus angulatus* McClure, 1995 (Decapoda: Caridea: Alpheidae). *Proceedings of the Biological Society of Washington* 115: 368-370.
- Melo, G.A.S. 2003. Famílias Atyidae, Palaemonidae e Sergestidae; p. 289-415 In G.A.S. Melo (ed.). *Manual de identificação dos Crustacea Decapoda de água doce do Brasil*. São Paulo: Edições Loyola.
- Milne-Edwards, A. 1878. Description de quelques espèces nouvelles de Crustacés provenant du voyage aux îles du Cap-Vert de MM. Bouvier et de Cessac. *Bulletin de la Société Philomathique de Paris* (7)2: 225-232.
- Milne Edwards, H. 1834-1840. Histoire naturelle des Crustacés, comprenant l'anatomie, la physiologie et la classification de ces animaux: 1-468, 1-532, 1-638, 1-32. Paris: Librairie encyclopédique de Roret.
- Neves, A., H.N. Cabral and L.S. Gordo. 2007. Distribution and abundance patterns of decapod crustaceans in the Sado estuary, Portugal. *Crustaceana* 80(1): 97-112.
- Ortmann, A. 1897. Os camarões da agua doce da America do Sul. *Revista do Museu Paulista* 2: 173-216.
- Rafinesque, C.S. 1815. *Analyse de la Nature, ou Tableau de l'Univers et des Corps Organisés*. Palermo: L'Imprimerie de Jean Barravecchia. 224 p.
- Ramos-Porto, M. 1980. Estudo ecológico da região de Itamaracá, Pernambuco, Brasil. VII. Crustáceos decápodos natantes. *Trabalhos Oceanográficos da Universidade Federal de Pernambuco* 15: 277-310.
- Ramos-Porto, M. 1986. Revisão das espécies do gênero *Leander* E. Desmarest que ocorrem no litoral brasileiro. *Trabalhos Oceanográficos da Universidade Federal de Pernambuco* 19: 7-26. [Dated 1985/86, published 1986].
- Ramos-Porto, M. & Coelho, P.A. 1990. Sinopse dos crustáceos decápodos brasileiros (família Palaemonidae). *Anais da Sociedade Nordestina de Zoologia* 3: 93-111.
- Rankin, W.M. 1898. The Northrop collection of Crustacea from the Bahamas. *Annals of the New York Academy of Sciences* 11: 225-258.
- Rathbun, M.J. 1900. Results of the Branner-Agassiz Expedition to Brazil. I. The decapod and stomatopod Crustacea. *Proceedings of the Washington Academy of Sciences* 2: 133-156.

- Rathbun, M.J. 1901. The Brachyura and Macrura of Porto Rico. *Bulletin of the United States Fish Commission* 20(2): 1-127.
- Rochebrune, A.T. de. 1883. Diagnoses d'Arthropodes nouveaux propres à la Sénégambie. *Bulletin de la Société Philomathique de Paris* (7)7: 167-182.
- Say, T. 1818. An account of the Crustacea of the United States, part 5. *Journal of the Academy of Natural Sciences at Philadelphia* 1: 235-253.
- Schmitt, W.L. 1924. Report on the Macrura, Anomura and Stomatopoda collected by the Barbados-Antigua Expedition from the University of Iowa in 1918. *University of Iowa Studies of Natural History* 10: 65-99, plates 1-5.
- Soledade, G.O. and A.O. Almeida. 2013. Snapping shrimps of the genus *Alpheus* Fabricius, 1798 from Brazil (Caridea: Alpheidae): updated checklist and key for identification. *Nauplius* 21(1): 89-122.
- Spence Bate, C.S. 1888. Report of the Crustacea Macrura collected by the H.M.S. "Challenger" during the years 1873-1876. *Report on the Scientific Results of the Voyage of H.M.S. "Challenger" During the Years 1873-1876*, *Zoology* 24: 1-942.
- Stimpson, W. 1871. Notes on North American Crustacea, in the museum of the Smithsonian Institution. No. III. *Annals of the Lyceum of Natural History in New York* 10: 119-163. [preprint issued in 1871, paper in bound volume in 1874 with page numbers 92-136]
- Udekem d'Acoz, C. d'. 1999. Inventaire et distribution des crustacés décapodes de l'Atlantique nord-oriental, de la Méditerranée et des eaux adjacentes au nord de 25°N. *Collection Patrimoines Naturels* (M.N.H.N./S.P.N.) 40 : 1-393p.
- Wicksten, M.K. and M.E. Hendrickx. 2003. An updated checklist of benthic marine and brackish water shrimps (Decapoda: Penaeoidea, Stenopodidea, Caridea) from the Eastern Tropical Pacific; p. 49-76 In M.E. Hendrickx (ed.). *Contributions to the Study of East Pacific Crustaceans* 2. [Contribuciones al Estudio de los Crustáceos del Pacífico Este 2]. México: Instituto de Ciencias del Mar y Limnología, UNAM.
- Wiegmann, A. 1836. Beschreibung einiger neuen Crustaceen des Berliner Museums aus Mexiko und Brasilien. *Archiv für Naturgeschichte* 2: 145-151.
- Williams, A.B. 1984. *Shrimps, Lobsters and Crabs of the Atlantic Coast of the Eastern United States, Maine to Florida*. Washington, D.C.: Smithsonian Institution Press. 550 p.

RECEIVED: August 2013

ACCEPTED: October 2013

PUBLISHED ONLINE: November 2013

EDITORIAL RESPONSIBILITY: Luis Ernesto Arruda Bezerra